

FAUNISTIC NOTE

The first record of *Boloria frigga* (Lepidoptera: Nymphalidae) on Kolguev Island, Arctic Russia

Grigory S. Potapov^{1,2}, Elizaveta A. Spitsyna¹, Vitaly M. Spitsyn¹

¹ N. Laverov Federal Center for Integrated Arctic Research of the Ural Branch of the Russian Academy of Sciences, Northern Dvina Emb. 23, Arkhangelsk, 163000, Russia

² Northern (Arctic) Federal University, Northern Dvina Emb. 17, Arkhangelsk, 163002, Russia

Corresponding author: Grigory S. Potapov (grigorij-potapov@yandex.ru)

Received 26 February 2021 | Accepted 11 November 2021 | Published 31 December 2021

Citation: Potapov GS, Spitsyna EA, Spitsyn VM (2021) The first record of *Boloria frigga* (Lepidoptera: Nymphalidae) on Kolguev Island, Arctic Russia. Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa" 64(2): 103–108. <https://doi.org/10.3897/travaux.64.e65131>

Abstract

A new record of *Boloria frigga* from the Arctic islands of European Russia is given. Image of a captured specimen, as well as brief considerations on the species habitat, are provided.

Keywords

Biogeography, Arctic islands, Barents Sea region, butterflies, fauna.

Frigga's fritillary, *Boloria frigga* (Thunberg, 1791), is a butterfly species with a circumboreal distribution. Its range includes Fennoscandia, the Baltic countries, Northern European Russia, Siberia, Mongolia, the Russian Far East, Alaska, the Rocky Mountains of the western states of the U.S., the Great Lakes, Quebec, and State of Maine (Tuzov and Bozano 2006; de Maynadier 2009; Haahtela et al. 2017; Söderström 2019).

In Fennoscandia and Northern European Russia, the range of *B. frigga* includes most of the part of this region, but it is a generally local species (Tatarinov 2016; Söderström 2019). Here, *B. frigga* prefers moss and shrub tundra, herb meadow patches in the tundra and raised bogs. Their caterpillars are known as the specialists and feed on the cloudberry (*Rubus chamaemorus*) (Tatarinov and Dolgin 1999). This is the main reason why *B. frigga* is rarer towards the south in the European part of

Russia, where the raised bogs are less common. The southern border of the species distribution is up to the Tver Oblast (Tuzov and Bozano 2006; Dubatolov et al. 2019). Here, in some regions of this part of European Russia, *B. frigga* is included in the Red List of Species, i.e. the Leningrad Oblast (2018), the Novgorod Oblast (2015), and the Tver Oblast (2016). The main threats to *B. frigga* are peat extraction and the draining of bogs.

The reliable records of *B. frigga* are relatively few in Northern European Russia. For example, there are no records of this species from the Arkhangelsk Oblast (Kozlov et al. 2014, 2017, 2020). The distribution of *B. frigga* is better acquainted for the Murmansk Oblast (Kozlov and Jalava 1994; Kozlov and Kullberg 2008), the Karelia Republic (Gorbach 2014), the Komi Republic, and the Nenets Autonomous Okrug (Bolotov 2011; Vlasova et al. 2014; Tatarinov 2016; Kozlov et al. 2019). Only two records of *B. frigga* are known on the Arctic islands of Northern European Russia, i.e. the Novaya Zemlya Archipelago and Dolgiy Island (review: Kullberg et al. 2019) (Table 1, Fig. 1).

Kolguev Island is one of the poorly studied territories of the European Arctic. This remote island has been little visited by entomologists. Only a few published information concerning the insect fauna, are known regarding this island. Six species of Nymphalidae have been recorded here previously, i.e. *Boloria chariclea* (Schneider, 1794), *B. polaris* (Boisduval, 1828), *Vanessa cardui* (Linnaeus, 1758), *Erebia disa* (Thunberg, 1791), *E. pandrose* (Borkhausen, 1788), *E. rossii* (Curtis, 1835) (Bolotov 2011; Kullberg et al. 2019). In all, no more than 21 species of Lepidoptera have been

Table 1. List of records of *Boloria frigga* in the north-eastern part of European Russia.

№	Locality (type)	N	E	References
1	Kolguev Island, 2 km N from Bugrino (settlement)	68°48'	49°17'	this study
2	Mesna (river)	–	–	Tatarinov 2016
3	Shoyna (settlement)	67°52'	44°09'	Tatarinov 2016
4	Shchuchye (lake)	67°26'	44°32'	Tatarinov 2016
5	Golodnaya Guba (lake)	67°56'	52°46'	Tatarinov 2016; Kozlov et al. 2019
6	Naryan-Mar (town)	67°38'	53°00'	Tatarinov 2016; Kozlov et al. 2019
7	Korovinskaya (bay)	68°11'	53°58'	Tatarinov 2016
8	Shapkina (river)	–	–	Tatarinov 2016
9	Dolgiy Island	69°16'	59°06'	Kullberg et al. 2019
10	Paiyakha (river)	69°20'	62°03'	Tatarinov 2016
11	Amderma (settlement)	69°45'	61°40'	Kozlov et al. 2019; Vlasova et al. 2014
12	Novaya Zemlya Archipelago	–	–	Kullberg et al. 2019

Notes: № corresponds to a number of the locality on the map (Fig. 1). With regards to several localities, it is impossible to determine the coordinates in a sufficiently accurate manner. These records are included without coordinates.




Figure 1. Map of the north-eastern part of European Russia. Records of *Boloria frigga* are shown as black dots, according the published data. The first record on Kolguev Island is presented as a red dot. Numbers for the localities are given in Table 1.

recorded so far from the Kolguev Island, making it one of the least studied islands from within Arctic islands of Russia (Kullberg et al. 2019; Spitsyn and Bolotov 2020). New records may allow us to create in the future a complete species list of Lepidoptera for this remote Arctic island. These data, regarding Lepidoptera in Arctic islands, are important in connection with the forecasts of changes in the ranges of species of insects under conditions of global warming.

A specimen of *B. frigga* was collected with an entomological net. It was prepared using the standard methods (Schauff 2001). The butterfly is deposited in the Russian Museum of Biodiversity Hotspots [RMBH], N. Laverov Federal Center for Integrated Arctic Research of the Ural Branch of the Russian Academy of Sciences (Arkhangelsk, Russia). The images of the specimen were taken with a Canon EOS 7D camera (Canon Inc., Tokyo, Japan). The source of the map is the online software SimpleMappr (Shorthouse 2010).

The first record: Russia, Kolguev Island (southern part), 2 km N from Bugrino near the Bugryanka River, moss-herb tundra, 68°48'04.1"N; 49°17'40.0"E, 16.vii.2020, Grigory Potapov leg. – 1♂ (Fig. 2).

B. frigga might be confused with *B. chariclea* (Schneider, 1794), *B. freija* (Thunberg, 1791), *B. polaris* (Boisduval, 1828). Upperside of wings of *B. frigga* with black heavy marking; upper hindwing with wide basal and discal black suffusion; under hindwing with a white basal spot at wing costa, white marginal spots absent (Tuzov and Bozano 2006).

B. frigga is widespread, but generally local in Northern European Russia. On the mainland, the number of records is quite a lot (Table 1, Fig. 1). Our record is located ca. 170 km northwards of the previous known locality of *B. frigga*. Before our study, we can indicate only two localities for the Arctic islands, where *B. frigga* was known.

Obviously, that further research in these remote tundra areas will increase the number of records of *B. frigga*. On Kolguev Island, the central and northern parts of this island with hills and river valleys are more prospective for the study. It seems that here, the Lepidoptera fauna is richer (Bolotov 2011).

The status of the population of *B. frigga* needs to be monitored in the European part of Russia. The disappearance of raised bogs, as a result of their economic use, has a negative impact on this species. The populations of *B. frigga* in the tundra zone appear more sustainable at the present time.

Acknowledgements

This study was supported by the Russian Ministry of Science and Higher Education (projects nos. 0409-2019-0042 to V.M.S. and 0409-2019-0039 to G.S.P), the Russian Foundation for Basic Research (project no. 19-34-90012) and the Young Scientists of Pomorye 2021 (project no. 07-2021a). Special thanks to A.G. Ardeev (Bugrino, Kolguev Island) for his invaluable help in material collecting. We are grateful to three anonymous reviewers for their valuable comments on earlier versions of this paper.




Figure 2. *Boloria frigga* on Kolguev Island: **A** General view of the specimen. Scale bar = 10 mm. (Photo: Elizaveta A. Spitsyna); **B** Habitat (moss-herb tundra) of this species on Kolguev Island (Photo: Grigory S. Potapov).

References:

- Bolotov IN (2011) Fauna and ecology of butterflies (Lepidoptera, Rhopalocera) from Kanin Peninsula and Kolguev Island. *Zoologicheskii zhurnal* 90(11): 1365–1373. [in Russian]
- Bublichenko YuN, Golubkov SM, Kiyashko PV (Chief Eds) (2018) The Red Book of the Leningrad Oblast. Animals. Papirus, Saint-Petersburg, 560 pp. [in Russian]
- De Maynadier PG (2009) *Boloria frigga saga* (Nymphalidae), a significant new record for Maine and Northeastern North America. *Journal of the Lepidopterists' Society* 63(3): 177–178.
- Dubatolov VV, Lvovskiy AL, Streltsov AN (2019) Nymphalidae. In: Sinev SYu (Ed.) Catalogue of the Lepidoptera of Russia. 2nd Edition. Zoological Institute of the RAS, Saint-Petersburg, 448 pp. [in Russian]
- Gorbach VV (2014) Fauna i ekologiya cheshuekrylyh (Lepidoptera: Hesperioidea et Papilionoidea) Vostochnoj Fennoscandii. Dr. Sc. thesis, Petrozavodsk State University, Petrozavodsk, Russia. [in Russian]
- Haahtela T, Saarinen K, Ojalainen P, Aarnio H (2017) Suomen ja Euroopan päiväperhoset. Gummerus, Helsinki, 383 pp. [in Finnish]
- Kozlov MV, Jalava J (1994) Lepidoptera of the Kola Peninsula, northwestern Russia. *Entomologica Fennica* 5: 65–85. <https://doi.org/10.33338/ef.83797>
- Kozlov MV, Kullberg J (2008) Lepidoptera of tundra habitats of the northern Kola Peninsula, Northwestern Russia. *Entomologica Fennica* 19: 114–121. <https://doi.org/10.33338/ef.84421>
- Kozlov MV, Kullberg J, Zverev VE (2014) Lepidoptera of Arkhangelsk oblast of Russia: a regional checklist. *Entomologica Fennica* 25: 113–141. <https://doi.org/10.33338/ef.48266>
- Kozlov MV, Kullberg J, Zverev VE (2017) New records of Lepidoptera from the Arkhangelsk oblast of Russia. *Entomologica Fennica* 28: 169–182. <https://doi.org/10.33338/ef.84685>
- Kozlov MV, Kullberg J, Zverev VE (2019) Moths and butterflies (Lepidoptera) of the continental part of the Nenets Autonomous Okrug, Russia. *Entomologica Fennica* 30: 72–89. <https://doi.org/10.33338/ef.82923>
- Kozlov MV, Kullberg J, Zverev VE (2020) Additions to the fauna of moths and butterflies (Lepidoptera) of the Arkhangelsk Oblast, Russia. *Annales Zoologici Fennici* 57: 183–194. <https://doi.org/10.5735/086.057.0119>
- Kullberg J, Filippov BYu, Spitsyn VM, Zubrij NA, Kozlov MV (2019) Moths and butterflies (Insecta: Lepidoptera) of the Russian Arctic islands in the Barents Sea. *Polar Biology* 42: 335–346. <https://doi.org/10.1007/s00300-018-2425-z>
- Orlov SV, Sokolov DL, Perova IS, Kokina OM, Turilov AV, Vedernikova ES (Eds) (2016) The Red Book of the Tver Oblast. Tverskoi pechatnyi dvor, Tver, 400 pp. [in Russian]
- Schauff ME (2001) Collecting and preserving insects and mites: techniques and tools. Systematic Entomology Laboratory, National Museum of Natural History, USDA, Washington, DC, 68 pp.
- Shorthouse DP (2010) SimpleMappr, an online tool to produce publication-quality point maps <https://www.simplemappr.net>
- Söderström B (2019) Nordens fjärilar – en fälthandbok. Bonnier Fakta, Stockholm, 272 pp. [in Swedish]

- Spitsyn VM, Bolotov NI (2020) First record of *Syngrapha diasema* (Lepidoptera: Noctuidae) from Kolguev Island, Arctic Russia. *Biharean Biologist* 14(1): 52–53.
- Tatarinov AG (2016) *Geografija dnevnyh cheshuekrylyh evropejskogo Severo-Vostoka Rossii*. KMK Press, Moscow, 255 pp. [in Russian]
- Tatarinov AG, Dolgin MM (1999) *Fauna evropejskogo Severo-Vostoka Rossii. Bulavousye cheshuekrylye*. Vol. 7, No. 1. Nauka, Saint-Petersburg, 183 pp. [in Russian]
- Tuzov VK, Bozano GC (2006) *Guide to the butterflies of the Palearctic region. Nymphalidae part II*. Omnes Artes, Milano, 72 pp.
- Vetkin YuE (Chief Ed) (2015) *The Red Book of the Novgorod Oblast*. Diton, Saint-Petersburg, 480 pp. [in Russian]
- Vlasova AA, Bolotov IN, Gofarov MY, Zubriy NA, Filippov BY, Frolov AA, Akimova IA (2014) Butterfly local faunas (Lepidoptera: Rhopalocera) of the European North of Russia: the north of Yugorsky Peninsula (Amderma) and Vaygach Island. *Vestnik Severnogo (Arkticheskogo) federalnogo universiteta. Seria: Estestvennye nauki* 3: 48–60. [in Russian]